

A scenario for the dynamics in the small entropy production limit

Leticia CUGLIANDOLO and Jorge KURCHAN¹

Laboratoire de Physique Théorique, École Normale Supérieure, 24 rue Lhomond, 75231 Paris Cedex 05, France

Laboratoire de Physique Théorique et Hautes Énergies, Jussieu, 4, Place Jussieu, 75005 Paris, France

¹*PMMH, École Supérieure de Physique et Chimie Industrielles, 10 rue Vauquelin, 75231 Paris Cedex 05, France*

(Received February 1, 2008)

We present a scenario for the nonequilibrium dynamics in the limit of small entropy production. We discuss (i) the appearance of different time-scales, (ii) the modification of the fluctuation-dissipation theorem and its relation to effective temperatures and partial equilibrations and (iii) the validity of Onsager reciprocity relations. We distinguish these properties by their reaction to infinitesimal perturbations. We recall that one can easily change the time dependence of observables by applying an infinitesimal force while time-reparametrization invariant features remain unchanged under the same perturbations. With the aim of better understanding these properties, we consider the effect of several baths with different temperatures and time-scales on the dynamics. This is done in two ways: numerically, by using a especially developed Monte Carlo algorithm that mimics the coupling to multiple baths; analytically, by computing the time-dependent probability density of simple systems in contact with multiple baths. We finally argue that these features are related to supersymmetry, the reparametrization invariance of the slow dynamics and its spontaneous breaking. This scenario is consistent within any perturbative resummation scheme. A brief version of this article appeared in Ref.¹⁾

KEYWORDS: nonequilibrium dynamics, effective temperatures

§1. Introduction

In many dynamical systems, there is a quantity that can be naturally interpreted as the entropy production.²⁾ It depends on the time-dependent probability distribution and vanishes only when the system reaches the Gibbs-Boltzmann (GB) measure. The limit of small, though non-vanishing, entropy production (SEP) is achieved in a relaxing system at long times, and in stationary systems driven by external forces (or by different thermal baths) when the external power input tends to zero.

A *finite* system in which all forces derive from a potential approaches, for any reasonable dynamics and in the long-time limit, the GB distribution. In contrast, an *extended* system might stay far from equilibrium even at very long times, producing entropy at a small rate. In this SEP limit the probability distribution *is not* a small perturbation around the GB equilibrium one.

There are several examples of such nonequilibrium systems. An infinitely large system with growing domains has, at any finite time, properties that are dominated by the domain walls³⁾ and is out of equilibrium. Similarly, glassy systems ‘age’: their responses and correlations depend not only on time-differences but also on the time elapsed since their preparation.^{4–11)} Both facts are taken into account neither by the GB distribution nor by a simple perturbation around it.

On the other hand, a system that would otherwise age might be kept in a stationary yet far from equilibrium regime of sustained (relative) youth by pumping a small quantity of energy.^{12–16)} Such thing would happen if domains are kept on average from growing by small, non-

conservative forces.¹⁴⁾ While the experimental situation for glasses is less clear in this respect, simple models suggest that if one subjects a system below the glass temperature to shearing forces, the viscosity may stop growing at a large, driving-power dependent value.¹⁷⁾ This situation is reminiscent of what happens in granular flow.¹⁸⁾ In these cases the systems are explicitly kept far from equilibrium by the ‘non-potential’ forces.

If the SEP limit is not the GB measure, is there any generic, model independent statement we can make about it? The purpose of this paper is to discuss several aspects of a scenario with partial equilibrations at separate time-scales¹⁾ inspired by ‘mean-field’ glass theory.¹⁰⁾ Although we cannot show that this scenario is the most general one, we can show that it is consistent in any dimension. It leads to definite predictions, some of which have already been tested. and some new ones that, to a limited extent, we test here.

The discussion will be aimed at extended systems satisfying reasonable spatial homogeneity properties. We may easily construct counterexamples of the present scenario in models with disconnected or almost disconnected parts, in which microscopic couplings scale with the system size, etc. Throughout this paper whenever a quantity is declared to be ‘small’, it must be understood with the zero entropy production limit taken first, and this itself taken after the infinite size limit.

§2. Formalism

Consider a system described by the variables ϕ_i , $i = 1, \dots, N$, that we encode in a vector $\vec{\phi}$, with energy $E(\vec{\phi})$. A nonconservative force $\vec{f}$, scaled with a parameter D , acts on $\vec{\phi}$. If the system is set in contact with a *white*

thermal bath, the uncorrelated Gaussian noises η_i have zero mean and variance $2\Gamma_0 T$. Γ_0 is the friction coefficient. The equations of motion are

$$-m_i \ddot{\phi}_i - \frac{\partial E(\vec{\phi})}{\partial \phi_i} + D f_i(\vec{\phi}) = \Gamma_0 \dot{\phi}_i - \eta_i \quad (2.1)$$

and the masses m_i can be zero in particular.

If the system is coupled to a *coloured* Gaussian bath,²⁰⁾ the noise and friction terms are

$$\int_{t_0}^t ds \nu(t, s) \dot{\phi}_i(s) - \rho_i \quad (2.2)$$

with $\langle \rho_i(t) \rho_j(t_w) \rangle = T^* \delta_{ij} \nu(t, t_w)$ and T^* the temperature of the bath. The presence of the same kernel $\nu(t, s)$ in the noise-noise correlation and in the friction term is a consequence of having taken a bath in equilibrium.

We define the correlation functions and the linear response of the variable ϕ_i to a kick applied to ϕ_j at t_w :

$$C_{ij}(t, t_w) = \langle \phi_i(t) \phi_j(t_w) \rangle, R_{ij}(t, t_w) = \left. \frac{\delta \langle \phi_i(t) \rangle}{\delta h_j(t_w)} \right|_{h=0}.$$

Using a standard diagrammatic approach,²¹⁾ one can derive a set of two coupled dynamic equations for $C_{ij}(t, t_w)$ and $R_{ij}(t, t_w)$ for *any* model. In general though, one does not succeed in performing the sum of diagrams involved. There are two main strategies to obtain a closed set of equations. The first consists in using simple models, i.e. simple expressions for $E(\vec{\phi})$. This is the choice made when working with $O(N)$ -type models for ferromagnetism,³⁾ fully-connected spin-glass models,^{10,22,23)} models in infinite dimensional embedding spaces,^{24,25)} etc. The second consists in choosing a recipe to select a subset of the infinite set of diagrams in such a way that one can sum them all and express this sum as an explicit functional of C_{ij} and R_{ij} . This is a route commonly followed in field theory and it is used, for example, to derive the mode-coupling theory for super-cooled liquids.^{10,26)}

In fully-connected models all n -point functions can be expressed in terms of two-point functions, which then provide a complete description of the dynamics. In finite dimensional models this cannot be done in closed, exact form, and the description in terms of two-point functions is not complete.

For all models, the equations for two-time functions have the structure of Schwinger-Dyson equations. Our claim is that in the SEP limit their solutions share the aspects discussed in Sect. 3. This is a consequence of the spontaneous breaking of a large reparametrization invariance appearing in the long waiting-time limit.¹⁾

We wish to stress that neither the methods used to obtain the dynamic equations nor the scenario discussed in this paper rely on the presence of quenched disorder. This is unnecessary in a dynamic treatment.

§3. The dynamic scenario

3.1 Fast and slow dynamics. Infinite sensitivity of the slow dynamics

Two-time functions have different behaviours in two time-regimes separated by a model-dependent characteristic time $\mathcal{T}(t_w)$ that diverges in the SEP limit.²⁷⁾ The re-

laxation is fast for $t - t_w$ small with respect to $\mathcal{T}(t_w)$ and it is fast for $t - t_w$ large with respect to $\mathcal{T}(t_w)$, $t - t_w \geq 0$. The correlation and response are then

$$C_{ij}(t, t_w) = C_{ij}^F(t - t_w) + \tilde{C}_{ij}(t, t_w), \\ R_{ij}(t, t_w) = R_{ij}^F(t - t_w) + \tilde{R}_{ij}(t, t_w),$$

where C_{ij}^F and R_{ij}^F are nonnegligible only for small time-differences, and $\tilde{C}_{ij}$ and $\tilde{R}_{ij}$ vary very slowly. This separation becomes sharper and sharper in the SEP limit. In this sense our treatment is asymptotic.

A very general feature of systems with slow dynamics is that *the time-dependence of $\tilde{C}_{ij}$ and $\tilde{R}_{ij}$ is sensitive to vanishingly small changes in the equations of motion*. The most familiar example is ferromagnetic coarsening in which an arbitrary small random field changes the growth law²⁸⁾ from power to log, making the slow part of the autocorrelation change from being a function of t_w/t to being a function of $\ln(t_w/\tau_0)/\ln(t/\tau_0)$. Another extreme example is the case of mean-field glasses, in which weak nonconservative forces completely destroy aging, rendering the problem stationary.¹³⁾

The “infinite sensitivity” is due to the presence of flat directions in the free-energy landscape^{27,29,30)} and this is also the very reason why there is slow dynamics. This physical fact has a mathematical counterpart in the invariances of the equations.¹⁾

The fact that one can change from aging to stationary slow dynamics by applying infinitesimal perturbations is indeed the justification for treating both situations on the same footing.

3.2 Fluctuation-dissipation theorem and natural effective temperatures: time-scales and thermalisation.

Quite generally, a weak form of the fluctuation-dissipation theorem (FDT) holds in the SEP limit³¹⁾

$$R_{ij}(t, t_w) - \beta \frac{\partial C_{ij}(t, t_w)}{\partial t_w} \Theta(t - t_w) \sim 0. \quad (3.1)$$

This is either because both terms are large and cancel leading to the usual FDT or because both are small but not necessarily equal, violating FDT. β is the inverse temperature of the thermal bath, itself in equilibrium. A stronger relation holds also in the SEP limit:

$$R_{ij}(t, t_w) = \beta_{ij}^{\text{EFF}}(t, t_w) \frac{\partial C_{ij}(t, t_w)}{\partial t_w} \Theta(t - t_w) \quad (3.2)$$

and defines the inverse ‘effective temperature’^{19,32–34)} β_{ij}^{EFF} . Equation (3.1) states that one can have effective temperatures that are different from $1/\beta$ only for the slow degrees of freedom. In what follows we call *natural effective temperatures* the values that T^{EFF} takes for a system in contact with a *single* bath, as opposed to those arising due to more complicated thermal environments. The existence of non-trivial effective temperatures is essentially related to the production of entropy.³¹⁾

If we define, as in experiments,⁵⁾

$$(\chi'_{ij} + i\chi''_{ij})(\omega, t_w) = \int_0^{t_w} d\tau e^{i\omega\tau} R_{ij}(t_w, t_w - \tau) \quad (3.3)$$

$$\tilde{C}_{ij}(\omega, t_w) = \int_0^{t_w} d\tau e^{i\omega\tau} C_{ij}(t_w, t_w - \tau) \quad (3.4)$$

β^{EFF} can be alternatively defined by using¹⁹⁾

$$\beta_{ij}^{\text{EFF}}(\omega, t_w) \equiv \frac{\chi_{ij}''(\omega, t_w)}{\omega \text{Re} \tilde{C}_{ij}(\omega, t_w)}. \quad (3.5)$$

Equations (3.2) and (3.5) do not necessarily coincide.

The scenario discussed in this paper states that *for all pairs of observables and for each time-scale the effective temperatures take a single value*. That is,

$$\beta_{ij}^{\text{EFF}}(\omega, t_w) \sim \beta^{\text{EFF}}(\omega, t_w), \forall i, j \quad (3.6)$$

$$\beta^{\text{EFF}}(\omega_1, t_w) - \beta^{\text{EFF}}(\omega_2, t_w) \sim 0, \text{ if } \frac{\omega_1}{\omega_2} = \text{cst} \quad (3.7)$$

in the SEP (old aging or weakly driven) limit. Importantly enough, the definitions (3.2) and (3.5) are in this limit completely equivalent. [Note, in passing, a peculiarity of the (important) case of infinite effective temperatures: all β 's in (3.6) and (3.7) might tend to zero but their quotient not be one.]

It is also important to mention that the value of T^{EFF} can sometimes very slowly change in time, in a much slower scale than all other in the problem, as emphasized by Nieuwenhuizen.³⁵⁾




Fig. 1. M copies of the system coupled to a heat bath at temperature T and to a thermometer that is, in this sketch, represented by an oscillator.

In order to show that $\beta^{\text{EFF}}(\omega_1, t_w)$ is indeed a temperature measured by a thermometer we proceed as in Ref.^{19,36)} We measure, with a ‘thermometer’ consisting of a degree of freedom x with a potential $V(x)$ (see Fig. 1), the temperature of an observable $A(\vec{\phi})$. In order to do so, we couple the thermometer linearly to the observable A of an ensemble of many (M) independent copies of the system, subject to the same history.³⁷⁾ In App. A we show that the equation of motion for x becomes, for large M ,

$$m\ddot{x}(t) = -\frac{\partial V(x)}{\partial x(t)} + k^2 \int_{t_0}^t ds R_{AA}(t, s)x(s) + \rho(t) \quad (3.8)$$

with ρ a Gaussian noise with zero mean and correlation

$$\langle \rho(t)\rho(t_w) \rangle = k^2 C_{AA}(t, t_w). \quad (3.9)$$

C_{AA} is the auto-correlation of the observable A at two different times and R_{AA} is the response of the same observable with respect to an infinitesimal perturbation h that acts at time t_w modifying the Hamiltonian according to $H \rightarrow H - A(t_w)h$. If R_{AA} and C_{AA} are linked by a time-scale dependent effective temperature that satisfies (3.7), we can decompose them as a sum of terms, each evolving in a different time-scale. Hence, in the weak coupling limit $k^2 \rightarrow 0$, Eq. (3.9) is a Langevin equation with several ‘baths’ of the form (2.2) acting on widely separated time-scales and each with its own temperature. If x is tuned to respond to a *single* time-scale, it will measure only its temperature.³⁸⁾

A simple way to view the relation of β^{EFF} to the zeroth law¹⁹⁾ is to connect alternatively x to an observable A and an observable B (simultaneously for an ensemble of M systems, as in the thermometer case). One shows that the average heat flow goes from the observable having a lower to the one having a higher β^{EFF} .

On the experimental side, Grigera and Israeloff³⁹⁾ gave evidence for the existence of non-trivial effective temperatures in glycerol by showing that Nyquist FDT is violated below T_g . In the context of spin-glasses, an indirect – and not exact – method also suggests that non-trivial effective temperatures appear in the glassy phase.⁴⁰⁾

On the numerical side, the simulations of many groups gave evidence for the existence of non-trivial FDT violations (and, consequently, effective temperatures) in a large variety of models. Among them we can mention models undergoing domain growth,⁴¹⁾ spin-glasses in finite dimensions,⁴²⁾ and Lennard-Jones systems.⁴³⁾

3.3 Response to slow, auxiliary baths. Thermalization of subsystems.

If glassy systems have natural effective temperatures associated with the slow degrees of freedom, it is then natural to ask how would they react to a small coupling to an additional slow auxiliary bath of temperature T^* .¹⁾ The effect of the additional bath is taken into account by adding a term like (2.2) to the Langevin equation. The simplest choice is $\nu(t, t_w) = \tilde{\nu}(|t - t_w|/\tau^*)$ with τ^* the characteristic time of the bath and $\tilde{\nu}$ a smooth decreasing function.

Within this scenario an arbitrarily weak auxiliary bath with $\tilde{\nu}(0)$ small has an important effect provided it is sufficiently slow (τ^* large), and the temperature T^* is within the range of effective temperatures of the slow dynamics. The (slow) time dependences of all correlation and response functions are affected by a time rescaling $t \rightarrow K(t)$ such that the time-scale which has a natural effective temperature T^* is locked to the time-scale τ^* . (This is possible thanks to the sensitivity property.)

In particular, an aging system with a single $T^{\text{EFF}} > T$ will become stationary if $T^* > T^{\text{EFF}}$, and be hardly affected if $T^* < T^{\text{EFF}}$. More generally, an aging system with multiple effective temperatures will become partially stationary (for all the time-scales with $T^* > T^{\text{EFF}}$),

but will still have aging for time-scales with $T^* < T^{\text{EFF}}$.

We hence have argued,¹⁾ very much in the spirit of Refs.^{45–48)} that the coupling to slow auxiliary baths are perturbations ‘conjugate to the natural temperatures’. The arguments put forward in the previous two paragraphs are supported by simulations of spin-glasses both mean-field and 3D (the latter very encouraging).¹⁾ In App. C we describe the algorithm used to simulate a spin system in contact with a multiple bath.

Another relevant question is mutual thermalisation between two subsystems having different natural effective temperatures, but sharing the same bath.¹⁹⁾ This is related to the response to an auxiliary slow bath, since, to a certain extent, each system acts as an auxiliary bath for the other when placed in mutual contact.

Within the present scenario two situations are possible:¹⁾ strong coupling, in which the effective temperatures equalise; and weak coupling, in which the combined system preserves essentially the temperatures of its constituents, but rearranges them in widely separated time-scales. In the latter case, the combined system has three temperatures, the bath temperature for the fast processes, the lowest and highest original effective temperatures for the intermediate and slowest timescales, respectively.

3.4 Reciprocity relations.

An unexpected feature that appears in this scenario are the reciprocity (Onsager) relations:¹⁾

$$\langle A(t)B(t_w) \rangle = \langle B(t)A(t_w) \rangle. \quad (3.10)$$

If this equality holds for the observables A and B , Eqs. (3.6) and (3.7) imply a similar relation for the responses. There is no reason *a priori* why the reciprocity relations should hold in a situation in which FDT is strongly violated. This is all the more surprising in an aging case in which the system is not even stationary. The interest of these relations is that they are relatively easy to measure in a simulation (results for 3DEA appeared in Ref.¹⁾) or in an experiment. The reciprocity relations are intimately related to the partial equilibrations.

§4. Multiply thermalized systems

A standard procedure in the dynamics of extended systems is to keep some macroscopic variables of interest and integrate away the others. The eliminated variables become part of a ‘thermal’ bath.

In *equilibrium*, the validity of FDT and of time-translational invariance (TTI) ensure that the friction and noise terms, coming from the projected sector, satisfy FDT (of the second kind). The whole procedure is self-consistent since the variables evolving in contact with a bath satisfying FDT will eventually satisfy FDT, now called of the first kind.²⁾

In an *out of equilibrium* situation FDT (and sometimes TTI) do not hold. How can we then picture the effect of the projected variables on our chosen ones?

We treat a system in contact with a multiple bath with different temperatures and time-scales that is based on

subsequent adiabatic approximations of the slower baths. This approach has points in common with the ‘pinning field’ approach of Kirkpatrick and Thirumalai⁴⁶⁾ and Monasson⁴⁷⁾ and with recent developments of Allahverdyan *et al.*⁴⁸⁾ and Franz and Virasoro⁴⁹⁾ We claim that, in the SEP limit, *correlation and response functions of the selected macroscopic variables behave as if they were subject to such a multiple bath.*

By studying in detail the evolution of a harmonic oscillator in contact with a multiple bath, we show how, even in this very simple model, different effective temperatures are induced by the coupling to several baths.⁵⁰⁾

4.1 The construction

Consider a particle of coordinate x moving in a potential $V(x)$ under the influence of two thermal baths: a fast white bath of temperature T and friction coefficient Γ_0 , and a slow bath of temperature T^* , characteristic time-scale τ^* and strength Γ_1 . Note that such a system has a non-Gibbsian stationary measure. We are interested in the evolution of the particle taking τ^* large with respect to any other characteristic time. (The following argument can be easily extended to any number of variables x_i , $i = 1, \dots, N$.) This problem, and its solution, is very similar to the one studied by Allahverdyan and Nieuwenhuizen in the second article of.⁴⁸⁾ These authors analysed two interacting variables that are respectively coupled to two baths with different temperatures and very different time-scales.

Neglecting the effect of inertia one has

$$\Gamma_0 \dot{x}(t) + \int_{-\infty}^t ds \, \tilde{\nu}(t-s) \dot{x}(s) = -\frac{\partial V}{\partial x(t)} + \eta(t) + \rho(t),$$

where η and ρ are the Gaussian thermal noises of the fast and slow baths, respectively, both with zero mean and variances $\langle \eta(t)\eta(t_w) \rangle = 2T\Gamma_0\delta(t-t_w)$ and $\langle \rho(t)\rho(t_w) \rangle = T^*\tilde{\nu}(|t-t_w|/\tau^*)$. We assume that $\tilde{\nu}(0) \equiv \Gamma_1$ and $\dot{\tilde{\nu}}(0^+) = 0$. We have set the system in contact with the slow bath at the initial time $t_0 = -\infty$. If we integrate by parts,

$$\Gamma_0 \dot{x}(t) = -\frac{\partial V}{\partial x(t)} - \Gamma_1 x(t) + \eta(t) + h(t) \quad (4.1)$$

$$h(t) \equiv -\int_{-\infty}^t ds \, \dot{\tilde{\nu}}(t-s)x(s) + \rho(t). \quad (4.2)$$

In the adiabatic limit, the slow bath generates a quasi-static field $h(t)$. (Indeed, using $\dot{\tilde{\nu}}(0^+) = 0$ and $\tau^* \gg 1$, one shows $\dot{h} = O(1/\tau^*) \ll 1$.) Hence, x has a fast evolution given by Eq. (4.1) with h fixed and it achieves a distribution

$$P(x/h) = \frac{e^{-\beta(V(x)+\Gamma_1 \frac{x^2}{2}-hx)}}{\int dx e^{-\beta(V(x)+\Gamma_1 \frac{x^2}{2}-hx)}}. \quad (4.3)$$

The denominator defines $Z(h)$ and $F(h) \equiv -\beta^{-1} \ln Z(h)$. Henceforth we denote $P(a/b)$ the conditional probability of a given b at stationarity.

The approximate evolution of h is given by Eq. (4.2) with the replacement of x in the friction term by its

average with respect to the fast evolution:

$$h(t) = \int_{-\infty}^t ds \dot{\nu}(t-s) \frac{\partial F(h)}{\partial h}(s) + \rho(t). \quad (4.4)$$

This equation is a non-Markovian and it has all the properties of a system coupled to a (slow) bath of temperature T^* . In particular, the stationary distribution is

$$\hat{P}(h) = \frac{e^{-\beta^*(F(h) + \frac{h^2}{2\Gamma_1})}}{\int dh e^{-\beta^*(F(h) + \frac{h^2}{2\Gamma_1})}}. \quad (4.5)$$

Reciprocity and FDT for functions of h also hold at stationarity. The stationary distribution of x now reads

$$P(x) = \int dh P(x/h) \hat{P}(h). \quad (4.6)$$

We prove these properties in App. B. In particular, if $T = T^*$ we recover the usual GB distribution for $P(x)$ regardless of the details of the slow bath.

Note that the distribution function $P(x)$ contains the superposition of different time-scales and is not a very eloquent quantity.

A case of great interest is the one of a slow bath that is not itself stationary, but ages. Suppose that we have

$$\nu(t, s) = \tilde{\nu}(L(s)/L(t)) \quad t \geq s \quad (4.7)$$

for some monotonically increasing function L . For the fast motion, at large enough times, we recover Eq. (4.3) while Eq. (4.2) follows from the change of variables

$$\mathcal{T} = \ln L(t), \quad \mathcal{T}' = \ln L(s). \quad (4.8)$$

In these new time-like variables, the rest of the derivation can be carried through identically. ‘Stationary’ means in this case invariant with respect to $L \rightarrow \Delta \times L$ that leads to $\mathcal{T} \rightarrow \mathcal{T} + \ln \Delta$ in time-like variables (Δ is a parameter).

4.2 Reciprocity and FDT

In its strongest form, reciprocity means that the joint probability of having x at time t and x' at t_w is equal *at stationarity*⁵¹⁾ to the probability of having x' at time t and x at t_w . To prove that this indeed happens, we use the separation of time-scales and consider two cases:

- i. If $t - t_w$ is small, h is approximately fixed and we have an equilibrium problem at temperature T , with a fixed field h . Reciprocity holds in the usual way.
- ii. If $t - t_w$ is such that $h(t) - h(t_w) \neq 0$, we have

$$P(x, t; x', t_w) = \int dh dh' P(x/h) P(h, t/h', t_w) P(h'/x') P(x') \quad (4.9)$$

Since $P(h, t; h', t_w) = P(h', t; h, t_w)$ (see App. B),

$$P(x, t; x', t_w) = P(x', t; x, t_w). \quad (4.9)$$

In a similar way one shows that FDT holds separately for each timescale with its own temperature.

- i. If $t - t_w$ is small FDT holds for each h . Denoting $\delta P(x, t)/\delta H_A(t_w)$ the variation of the distribution at time t due to a field conjugate to A that has been on from $-\infty$ to t_w , we have

$$\frac{\delta P(x, t)}{\delta H_A(t_w)} = \beta \int dx' P(x, t/x', t_w) A(x') P(x'). \quad (4.10)$$

The integrated form of FDT for two observables A and B can be obtained by multiplying by $B(x)$ and integrating over x .

- ii. If $t - t_w$ is such that $h(t) - h(t_w) \neq 0$, while the field conjugate to A is on, h evolves with Eq. (4.4) and

$$\begin{aligned} F_A(h) &= -\frac{1}{\beta} \ln \int dx' e^{-\beta(V(x') - H_A A(x') + \Gamma_1 \frac{x'^2}{2} - hx')} \\ &= F(h) - H_A A(h), \end{aligned} \quad (4.11)$$

with $\mathcal{A}(h) = \int dx' P(x'/h) A(x')$. If H_A is turned off at t_w , $P(x/h)$, is first modified in a short time-scale with respect to that of h . Then, h continues to evolve but with a modified effective potential $F(h)$. Now, FDT holds for the evolution of h at temperature T^*

$$\frac{\delta \hat{P}(h, t)}{\delta H_A(t_w)} = \beta^* \int dh' P(h, t/h', t_w) \mathcal{A}(h') \hat{P}(h'),$$

(see App. B) and

$$\begin{aligned} \frac{\delta P(x, t)}{\delta H_A(t_w)} &= \int dh P(x/h) \frac{\delta \hat{P}(h, t)}{\delta H_A(t_w)} \\ &= \beta^* \int dh dh' P(x/h) P(h, t/h', t_w) \mathcal{A}(h') \hat{P}(h'). \end{aligned}$$

Using $\hat{P}(h') \mathcal{A}(h') = \int dx' P(x') P(h'/x') A(x')$ we obtain

$$\frac{\delta P(x, t)}{\delta H_A(t_w)} = \beta^* \int dx' P(x, t/x', t_w) A(x') P(x').$$

For separate times FDT holds with temperature T^* .

4.3 The effect of a weak coloured bath

The simplest example where to check the ideas described in Sect. 4.1 and 4.2 is a harmonic oscillator in contact with a white and a coloured bath with exponential correlation. This problem can be tackled using a variety of techniques.

From the exact asymptotic solution we extract the behaviour of the integrated response $\chi(\tau) \equiv \int_0^\tau d\tau' R(\tau')$ vs. $C(\tau)$ and read T^{EFF} from it. In Fig. 2 we show $\chi(C)$ for $T = 0.5$, $\Gamma_0 = 1$, $T^* = 1$. In the plot above, $(\omega + \Gamma_1)\tau^* = 2000 \gg \Gamma_0 = 1$, with ω the frequency of the oscillator. The evolution takes place in two time-scales (or correlation-scales) characterised by temperatures T and T^* . The straight lines have slopes $-1/T$ and $-1/T^*$ showing that there are two temperatures associated to the motion of the particle: a fast motion for $q < C < q_d$ controlled by T and a slow motion for $0 < C < q$ that is instead controlled by T^* . In the plot below $(\omega + \Gamma_1)\tau^* = 2 = O(\Gamma_0 = 1)$, the time-scales (or correlation-scales) are not well separated, and $\chi(C)$ continuously interpolates between a region of slope of $-1/T$ and a region of slope of $-1/T^*$.

This problem can be alternatively analysed using the technique described in Sect. 4.1. One recovers, as expected, the exact results of the previous paragraphs.

4.4 Many scales

The construction of Sect. 4.1 can be generalised to many nested scales. Let us see how this is done for three




Fig. 2. The $\chi(C)$ plot for a harmonic oscillator subject to a white bath of temperature $T = 0.5$ and a coloured noise of temperature $T^* = 1$. Above: $(\omega + \Gamma_1)\tau^* = 2000 \gg \Gamma_0 = 1$. The straight lines (dots) have inverse slopes $-1/T = -2$ and $-1/T^* = -1$. Below: $(\omega + \Gamma_1)\tau^* = 2 = O(\Gamma_0 = 1)$.

baths; the equation of motion is

$$\Gamma_0 \dot{x}(t) + \int_{-\infty}^t ds (\nu_1(t-s) + \nu_2(t-s)) \dot{x}(s) = -\frac{\partial V(x)}{\partial x(t)} + \eta(t) + \rho_1(t) + \rho_2(t) \quad (4.12)$$

with ρ_1 and ρ_2 two Gaussian thermal noises with zero mean, $\langle \rho_1(t) \rho_1(t_w) \rangle = T_1^* \tilde{\nu}_1(|t - t_w|/\tau_1^*)$, $\langle \rho_2(t) \rho_2(t_w) \rangle = T_2^* \tilde{\nu}_2(|t - t_w|/\tau_2^*)$ and $1 \ll \tau_1^* \ll \tau_2^*$. Proceeding as before, we write

$$\Gamma_0 \dot{x}(t) = -\frac{\partial V(x)}{\partial x(t)} + (\Gamma_1 + \Gamma_2) x(t) + \eta(t) + h_1(t) + h_2(t)$$

with $h_i(t) \equiv -\int_{-\infty}^t ds \dot{\nu}_i(t-s)x(s) + \rho_i(t)$, $i = 1, 2$, and $\nu_i(0) = \Gamma_i$, $\dot{\nu}_i(0^+) = 0$, $i = 1, 2$.

In the fastest evolution x achieves distribution,

$$P(x/h_1, h_2) = \frac{e^{-\beta(V(x) + (\Gamma_1 + \Gamma_2)\frac{x^2}{2} - (h_1 + h_2)x)}}{Z_{h_1, h_2}}, \quad (4.13)$$

with Z_{h_1, h_2} ensuring $\int dx P(x/h_1, h_2) = 1$. The evolution of h_1 is obtained by taking h_2 to be adiabatic

$$h_1(t) = \int_{-\infty}^t ds \dot{\nu}_1(t-s) \frac{\partial F_1(h_1 + h_2)}{\partial h_1}(s) + \rho_1(t) \quad (4.14)$$

with $e^{-\beta^* F_1(h_1 + h_2)} \equiv \int dx e^{-\beta(V(x) + (\Gamma_1 + \Gamma_2)\frac{x^2}{2} - (h_1 + h_2)x)}$ yielding a stationary distribution for h_1

$$P(h_1/h_2) \propto \exp \left[-\beta_1^* \left(F_1(h_1 + h_2) + \frac{h_1^2}{2\Gamma_1^2} \right) \right]. \quad (4.15)$$

Finally, h_2 evolves at inverse temperature β_2^* and has a

stationary distribution:

$$\hat{P}(h_2) \propto \exp \left[-\beta_2^* \left(F_2(h_2) + \frac{h_2^2}{2\Gamma_2^2} \right) \right] \quad (4.16)$$

with $e^{-\beta_2^* F_2(h_2)} \equiv \int dh_1 e^{-\beta_1^* \left(F_1(h_1 + h_2) + \frac{h_1^2}{2\Gamma_1^2} \right)}$. One can also consider two slow baths that are non-stationary. In order to have two well separated time-scales, we need two (increasing) scaling functions $L_1(t)$, $L_2(t)$ with

$$\lim_{t \rightarrow \infty} (d_t \ln L_2(t) / d_t \ln L_1(t)) = 0 \quad (4.17)$$

and, for example,

$$\nu_1(t, s) = \tilde{\nu}(L_1(s)/L_1(t)), \quad \nu_2(t, s) = \tilde{\nu}(L_2(s)/L_2(t)),$$

$t \geq s$. After performing two transformations like the ones in Eq. (4.8), the rest of the derivation can be repeated.

For this example one concludes that the ‘probability cloud’ of each scale acts as a configuration for the slower scale — moving in an effective potential.

§5. Conclusions

In the last years great progress has been made in understanding the asymptotic nonequilibrium evolution of systems with slow dynamics. Much of these developments came from the study of simplified models that include classical glassy disordered and non-disordered models, quantum disordered systems and, importantly enough, systems that are constantly driven out of equilibrium by external forces. The results collected from the solution of several of these cases made apparent the existence of various general features. In this article (as well as in Ref.¹⁾) we stressed which are the features that we believe build a scenario for the slow dynamics in the limit of small entropy production. Properties *ii-iv* discussed in Sect. 3. can be quite surprising, in particular the latter. In addition, in view of the infinite sensitivity of the slow time dependence of the correlation and response functions, it is a non-trivial fact that properties *ii-iv* are preserved under small perturbations. In Ref.¹⁾ we justified this fact by studying the invariances (and breakdown of) the dynamic equations in the SEP limit.⁵⁴⁾

In short, the image we have in mind is better grasped by comparing the general case to the simple problem studied in Sect. 4. The Langevin equation for a variable x including a non-local friction kernel and a correlated noise, Eq. (4.1), is reminiscent of the equation for a single effective variable that one obtains in fully connected models, large dimension approximations, etc.^{25, 52)} Indeed, in the large N limit, one can derive such an equation by performing a saddle-point approximation of the dynamic generating functional. For any dynamic variable ϕ_i , the ‘single-variable equation’ reads

$$\Gamma_0 \dot{\phi}_i(t) = -\mu(t) \phi_i(t) + \int_{t_0}^t ds \Sigma(t, s) \phi_i(s) + \rho_i(t) + \eta_i(t). \quad (5.1)$$

There are two noise sources in this equation: $\eta_i(t)$ is the original white noise while $\rho_i(t)$ is an effective (Gaussian) noise with zero mean and correlations self-consistently given by $\langle \rho_i(t) \rho_j(t_w) \rangle = \delta_{ij} T^* D(t, t_w)$. The

vertex $D(t, t_w)$ plays the rôle of the coloured noise correlation in a usual Langevin equation. The self-energy $\Sigma(t, t_w)$ appears here as an ‘integrated friction’.

In the case of a system in contact with an external coloured bath, the correlation of the noise $T^* \nu(t, t_w)$ is simply related to the retarded friction $\nu(t, t_w)$. This is the statement of FDT applies for the bath. In the case of the single-variable equation for a more complicated model, the friction kernel $\Sigma(t, t_w)$ and the correlation of the coloured noise $T^* D(t, t_w)$ are properties of the system and are not necessarily related in a simple manner. In the SEP limit, they get related by a modification of FDT with time-scale dependent effective temperature $T^{\text{EFF}}(t, t_w)$.

One then concludes that the structure of these two problems is indeed very similar:

- If one weakly couples a simple system to a slow bath of temperature T^* , at sufficiently slow time scales the system acquires the temperature T^* .
- Glassy systems arrange their internal degrees of freedom in such a way that slow degrees of freedom select their own effective temperatures.

Implicit in the construction presented in Section 4.1 is the fact that all observables have the same effective temperatures at the same two-times. Also implicit are the reciprocity relations. These are important in that they are potentially measurable, and quite unexpected out of equilibrium.

Appendix A: Measurements of natural effective temperatures

We here recall the definition of a time-scale dependent effective temperature, for systems out of equilibrium, given in Ref.¹⁹⁾ The presentation follows closely this reference but it makes some points of the derivation more precise.

Let us consider M non-interacting copies of the system,³⁷⁾ and couple each of them to a simple system that acts as a thermometer in the manner sketched in Fig. 1. For simplicity, we describe this thermometer with a single variable x and each system with a variable $\vec{\phi}_\alpha$, $\alpha = 1, \dots, M$. The total energy of the complex is

$$E_{\text{TOT}} = m \frac{\dot{x}^2}{2} + V(x) + \sum_{\alpha=1}^M E(\vec{\phi}_\alpha) - \frac{k}{M^{1/2}} x \sum_{\alpha=1}^M A(\vec{\phi}_\alpha),$$

where $V(x)$ and $E(\vec{\phi}_\alpha)$ are the potential energies of the isolated thermometer and each isolated system. For each copy the last term yields an infinitesimal (for M large) perturbation corresponding to a field $kx/\sqrt{M}$ conjugate to the observable $A(\vec{\phi}_\alpha)$. The equation of motion for x is

$$m\ddot{x}(t) = -\frac{\partial V(x)}{\partial x(t)} - \frac{k}{M^{1/2}} \sum_{\alpha=1}^M A(\vec{\phi}_\alpha)(t). \quad (\text{A.1})$$

For simplicity we choose an operator A such that $\langle A(\vec{\phi}_\alpha) \rangle_{k=0} = 0$ where $\langle \bullet \rangle$ represents either an average over different histories of the same system or an average over different systems, e.g. $\langle f \rangle = 1/M \sum_{\alpha=1}^M f_\alpha$. (If the average of the operator A does not vanish, one has to

work with the difference between the operator and its average.) The subindex $k = 0$ indicates that the average is taken in the absence of the external field $kx/M^{1/2}$. Whenever we take the average in the presence of the field we shall denote it $\langle \bullet \rangle_k$.

Assuming that linear response holds for each system, the total variation of the average $\langle A(\vec{\phi}_\alpha) \rangle_k$ caused by a field that has been applied from $t = 0$ upto t reads

$$\delta \langle A(\vec{\phi}_\alpha) \rangle_k(t) = \langle A(\vec{\phi}_\alpha) \rangle_k(t) - \langle A(\vec{\phi}_\alpha) \rangle_{k=0}(t) = \langle A(\vec{\phi}_\alpha) \rangle_k(t) - \int_0^t ds R_{A_\alpha, A_\alpha}(t, s) \frac{k}{\sqrt{M}} x(s), \quad (\text{A.2})$$

where $R_{A_\alpha, A_\alpha}(t, s)$ is the linear response of the observable $A(\vec{\phi}_\alpha)$ at time t to a change in energy $-k/\sqrt{M} x A(\vec{\phi}_\alpha)$ at time s . These thermal history-averaged responses are equal for all systems, we henceforth denote them $R(t, s)$ for all α . (We also simplify the notation by eliminating the subindex A that identifies the observable.) Adding the last equation over α and multiplying by $k/\sqrt{M}$ we have

$$\frac{k}{\sqrt{M}} \sum_{\alpha=1}^M \langle A(\vec{\phi}_\alpha) \rangle_k(t) = k^2 \int_0^t ds R(t, s) x(s). \quad (\text{A.3})$$

Adding and subtracting from the rhs of Eq. (A.1) the average (A.3), we recast it as

$$m\ddot{x}(t) = -\frac{\partial V(x)}{\partial x(t)} + k^2 \int_0^t ds R(t, s) x(s) + \rho(t) \quad (\text{A.4})$$

with

$$\rho(t) \equiv \frac{k}{\sqrt{M}} \sum_{\alpha=1}^M \left(A(\vec{\phi}_\alpha)(t) - \langle A(\vec{\phi}_\alpha) \rangle_k(t) \right). \quad (\text{A.5})$$

The ‘force’ $\rho(t)$ is the sum of M independent identically distributed random variables. For large M it becomes a Gaussian variable with zero mean and variance:

$$\langle \rho(t) \rho(t_w) \rangle_k = \frac{k^2}{M} \sum_{\alpha, \beta=1}^M C_{\alpha\beta}^{\text{CONN}}(t, t_w) \Big|_k. \quad (\text{A.6})$$

with $C_{\alpha\beta}^{\text{CONN}}(t, t_w) \Big|_k = \langle (A(\vec{\phi}_\alpha)(t) - \langle A(\vec{\phi}_\alpha) \rangle_k(t)) \times (A(\vec{\phi}_\beta)(t_w) - \langle A(\vec{\phi}_\beta) \rangle_k(t_w)) \rangle_k$ the connected correlation in the presence of the field. Up to leading order in k

$$C_{\alpha\beta}^{\text{CONN}}(t, t_w) \Big|_k \sim C_{\alpha\beta}^{\text{CONN}}(t, t_w) \Big|_{k=0}. \quad (\text{A.7})$$

The connected correlation can also be substituted by the usual correlation since $A(\vec{\phi}_\alpha)(t) = O(1)$ implies $\langle A(\vec{\phi}_\alpha)(t) \rangle_k = O(1/\sqrt{M})$. Thus

$$C_{\alpha\beta}^{\text{CONN}}(t, t_w) \Big|_{k=0} \sim C_{\alpha\beta}(t, t_w) \Big|_{k=0} \sim \delta_{\alpha\beta} C(t, t_w) \quad (\text{A.8})$$

since the bare systems are completely independent. Finally,

$$\langle \rho(t) \rho(t_w) \rangle_k = k^2 C(t, t_w). \quad (\text{A.9})$$

Thus the dynamic equation governing the evolution of the thermometer reads

$$m\ddot{x}(t) = -\frac{\partial V(x)}{\partial x(t)} + \rho(t) + k^2 \int_0^t ds R(t, s) x(s),$$

$$\langle \rho(t) \rho(t_w) \rangle_k = k^2 C(t, t_w). \quad (\text{A}\cdot 10)$$

If the response and the correlation of the system are related through the FDT relation

$$R(t, s) = \frac{1}{T} \frac{\partial C(t, s)}{\partial s} \Theta(t - s). \quad (\text{A}\cdot 11)$$

one integrates by parts the last term on the rhs of Eq. (A·10) and obtains

$$m\ddot{x}(t) = -\frac{\partial V}{\partial x(t)} + \rho(t) - \frac{k^2}{T} \int_0^t ds C(t, s) \dot{x}(s) + \frac{k^2}{T} (C(t, t)x(t) - C(t, 0)x(0)). \quad (\text{A}\cdot 12)$$

that in the limit of weak coupling, $k^2 \rightarrow 0$, becomes a Langevin equation since the last term can be neglected.

From this equation we can conclude that the thermometer, being a system with fast dynamics, will eventually equilibrate at the temperature that relates response and correlation of the system, Eq. (A·11). *The system acts as a thermal bath on the thermometer.*

We can also derive this result by choosing a harmonic oscillator as a thermometer.¹⁹⁾ Using $V(x) = m\omega_o^2 x^2/2$, one easily shows that the averaged potential energy of the thermometer over a time-window around the measuring time t_w is

$$\frac{1}{2} \langle E_{\text{osc}} \rangle_{t_w} = \frac{1}{2} \omega_o^2 \langle x^2 \rangle_{t_w}, \quad (\text{A}\cdot 13)$$

with ω_o is the probing frequency of the oscillator and $m = 1$. In the large t_w limit, it reaches the following limit

$$\langle E_{\text{osc}} \rangle_{t_w} = \frac{\omega_o \tilde{C}(\omega_o, t_w)}{\chi''(\omega_o, t_w)} \quad (\text{A}\cdot 14)$$

with $\tilde{C}$ and χ'' defined in Eqs. (3.3) and (3.4). If equipartition holds for the oscillator, one has

$$T^{\text{EFF}}(\omega_o, t_w) = \frac{\omega_o \tilde{C}(\omega_o, t_w)}{\chi''(\omega_o, t_w)}, \quad (\text{A}\cdot 15)$$

i.e. the definition given in Eq. (3.5).

Under the assumption (3.6) we can decompose the correlations and responses as follows:

$$C(t, t_w) = C^F(t, t_w) + C^1(t, t_w) + C^2(t, t_w) + \dots$$

$$R(t, t_w) = R^F(t, t_w) + R^1(t, t_w) + R^2(t, t_w) + \dots$$

with

$$R^i(t, t_w) = \beta^i \frac{\partial C^i(t, t_w)}{\partial t_w} \Theta(t - t_w), \quad (\text{A}\cdot 16)$$

$i = 0, 1, \dots$, $i = 0$ corresponding to the FDT scale. Then, Eqs. (3.9) and (A·12) corresponds to a system coupled to a series of baths of inverse temperatures $\beta^F, \beta^1, \beta^2$ acting on widely separated scales. The probing frequency of the thermometer can be selected to measure any of the values of the effective temperature.³⁸⁾

Actually, the effective temperature T^{EFF} in Eq. (A·15) might depend on the observable A considered. The picture in this paper describes the way in which we think these dependences arrange in a physical system.

Appendix B: Stationary distribution of the quasi-static field

We show that Eq. (4.4) leads to Eq. (4.5) and that at stationarity reciprocity and FDT hold. The Fourier transform of Eq. (4.4) reads

$$h(\omega) = r(\omega) \frac{\partial F}{\partial h}(\omega) + \rho(\omega), \quad (\text{B}\cdot 1)$$

$$\langle \rho(\omega) \rho(\omega') \rangle = T^* \nu(\omega) 2\pi \delta(\omega + \omega'), \quad (\text{B}\cdot 2)$$

where $r(\omega) \equiv \int_{-\infty}^{\infty} d\tau \exp(i\omega\tau) \dot{\nu}(\tau) \theta(\tau)$ is the slow bath's response function. It is then analytical in the upper half complex plane and satisfies, for real ω ,⁵³⁾ $r^*(\omega) = r(-\omega)$. FDT holds for the bath: $r(\omega) - r(-\omega) = -i\omega\nu(\omega)$. In order to prove Eq. (4.5), we introduce a set of auxiliary variables y_j satisfying the ordinary Langevin equation:

$$\left[m_j \frac{d^2}{dt^2} + \Gamma_j \frac{d}{dt} + \Omega_j \right] y_j = \xi_j(t) - \frac{\partial F(\sum_j A_j y_j)}{\partial y_j} \quad (\text{B}\cdot 3)$$

with $\langle \xi_i(t) \xi_j(t_w) \rangle = 2T^* \Gamma_j \delta_{ij} \delta(t - t_w)$. The aim is to show that $\sum_j A_j y_j$, satisfies the same equations as h , Eqs. (B·1) and (B·2).⁵⁵⁾ The introduction of y_i allows us to convert the original non-Markovian problem into a Markovian one. If F is such that Eq. (B·3) drives the ensemble y_i to equilibrium,

$$P(\vec{y}) \propto e^{-\beta^* \left(\frac{\Omega_j y_j^2}{2} + F(\sum_j A_j y_j) \right)}. \quad (\text{B}\cdot 4)$$

In Fourier space, Eq. (B·3) reads

$$[-m_j(\omega - \omega_j^+)(\omega - \omega_j^-) + \frac{\partial F(\sum_j A_j y_j)}{\partial y_j}] y_j(\omega) = \xi_j(\omega)$$

where $\omega_j^\pm$ are the roots of $-m_j \omega^\pm{}^2 - i\omega^\pm \Gamma_j + \Omega_j = 0$. Defining $h(\omega) = \sum_j A_j y_j(\omega)$ one has

$$h(\omega) = - \sum_j \frac{A_j \xi_j(\omega) + A_j^2 \partial_h F(h)}{m_j(\omega - \omega_j^+)(\omega - \omega_j^-)}. \quad (\text{B}\cdot 5)$$

Since $\text{Im}(\omega_j^\pm) \leq 0$, we can choose the variables y_j and the parameters m_j, Γ_j, Ω_j in such a way to identify

$$r(\omega) = \sum_j \frac{A_j^2}{m_j(\omega - \omega_j^+)(\omega - \omega_j^-)}. \quad (\text{B}\cdot 6)$$

Furthermore, we associate the first term in the rhs of Eq. (B·5) to the coloured noise $\rho(\omega)$ and we use Eq. (B·3) to obtain the noise-noise correlation:

$$\begin{aligned} \langle \rho(\omega) \rho(\omega') \rangle &= 2T^* \delta(\omega + \omega') \\ &\times \sum_j \frac{\Gamma_j A_j^2}{m_j^2(\omega - \omega_j^+)(\omega - \omega_j^-)(\omega + \omega_j^+)(\omega + \omega_j^-)} \end{aligned} \quad (\text{B}\cdot 7)$$

The properties of the poles $(\omega_j^\pm)^* = -\omega_j^\mp$ and $m_j(\omega_j^+ + \omega_j^-) = -i\Gamma_j$ yield FDT $\nu(\omega) = -\frac{1}{i\omega} [r(\omega) - r(-\omega)]$.

We need an expression for $\Gamma_1 \equiv \nu(t = 0) = \int_{-\infty}^{\infty} d\omega / (2\pi) \nu(\omega)$. Completing the integration over the upper or the lower half plane, we have

$$\Gamma_1 = \sum_j \frac{A_j^2}{\Omega_j} \quad (\text{B}\cdot 8)$$

We have shown that Eq. (4.4) is equivalent to an ordinary Markovian equation for $\vec{y}$, provided one identifies $h = \sum A_j y_j$. The stationary distribution for the y_j is then given by Eq. (B.4) $\hat{P}(h)$ is calculated by introducing a delta-function:

$$\begin{aligned} \hat{P}(h) &\propto \int d\lambda \int \Pi_j dy_j e^{-\beta^* \left(\frac{\Omega_j y_j^2}{2} + F(h) \right) - i\beta^* \lambda (\sum A_j y_j - h)} \\ &\propto \int d\lambda e^{-\beta^* \left(\sum \frac{A_j^2}{2\Omega_j} \lambda^2 - i\lambda h + F(h) \right)} \\ &\propto e^{-\beta^* \left(\frac{h^2}{2\Gamma_1} + F(h) \right)}. \end{aligned} \quad (\text{B.9})$$

The reciprocity property is immediate from the fact that, for a Markovian Langevin process in equilibrium,

$$P(\vec{y}, t; \vec{y}', t_w) = P(\vec{y}', t; \vec{y}, t_w). \quad (\text{B.10})$$

Similarly FDT follows from the fact that because it holds for any two functions of the y_j , it holds in particular for any two functions of h .

Appendix C: A Montecarlo algorithm with several baths

We describe the algorithm used to simulate the evolution of a system interacting with several thermal baths of different type. Let us consider a dynamic spin $s = \pm 1$ with energy $E(s)$ that is in contact with a ‘fast’ bath of inverse temperature β and a ‘slow’ bath of inverse temperature β^* and time-scale τ^* . The generalization to several variables and many baths is straightforward.

The algorithm is as follows.

i With probability $1/\tau^*$ we generate a Gaussian random variable h with s -dependent mean and variance:

$$\langle h \rangle = \beta^* \Gamma_1^2 s, \quad \langle h^2 \rangle - \langle h \rangle^2 = \Gamma_1^2. \quad (\text{C.1})$$

Γ_1 is the strength of the bath.

ii With probability $1 - 1/\tau^*$ we make ordinary Montecarlo updates of s at temperature $1/\beta$, with an energy $E_{\text{TOT}} = E(s) - hs$.

We justify this algorithm as follows. We model the slow bath by a system of $\mathcal{N}$ non-interacting spins $\sigma_a = \pm 1$, $a = 1, \dots, \mathcal{N}$, weakly coupled to s . The total energy of system and bath is given by

$$E_{\text{TOT}} = E(s) - \frac{\Gamma_1}{\mathcal{N}^{1/2}} s \sum_a \sigma_a = E(s) - hs, \quad (\text{C.2})$$

with $E(s)$ the energy of the free spin, Γ_1 the coupling constant and the ‘field’ h defined as $h \equiv \Gamma_1/\mathcal{N}^{1/2} \sum_a \sigma_a$.

We propose the following dynamics to the coupled system:

i Frequent updates of s keeping σ_a fixed. The spin is then in presence of a constant external field h . These updates are done with any Montecarlo procedure, at inverse temperature β .

ii Unfrequent updates of all σ_a , i.e. of the field h , keeping s fixed. This evolution is controlled by an inverse temperature β^* .

If the sequence is long, one can assume that each σ_a is in thermal equilibrium, at temperature $1/\beta^*$ with an energy $E(\sigma_a) \sim -\Gamma_1/\mathcal{N}^{1/2} s \sigma_a$. Each configuration $\sigma_a =$

± 1 has probabilities p_+ and p_- given by:

$$p_{\pm} = \frac{\exp(\pm \beta^* \frac{\Gamma_1}{\mathcal{N}^{1/2}} s)}{2 \cosh(\beta^* \frac{\Gamma_1}{\mathcal{N}^{1/2}} s)}. \quad (\text{C.3})$$

The σ_a are then identically distributed independent random variables. It follows that, in the limit of large $\mathcal{N}$, h is a Gaussian random variable. Its mean and variance are given by:

$$\begin{aligned} \langle h \rangle &= \frac{\Gamma_1}{\mathcal{N}^{1/2}} \sum_a \langle \sigma_a \rangle = \beta^* \Gamma_1^2 s + O\left(\frac{1}{\mathcal{N}^{1/2}}\right) \\ \langle h^2 \rangle - \langle h \rangle^2 &= \frac{\Gamma_1^2}{\mathcal{N}} \sum_a (\langle \sigma_a^2 \rangle - \langle \sigma_a \rangle^2) = \Gamma_1^2 + O\left(\frac{1}{\mathcal{N}^{1/2}}\right) \end{aligned}$$

As a check of the consistency of this procedure we examine if detailed balance holds, as expected, in the particular case of a variable coupled to two baths with equal temperatures $\beta = \beta^*$. The transition probability per unit time to go from the state s_A to the state s_B is

$$P(A \rightarrow B) = \frac{\Gamma_1}{\sqrt{2\pi}} \int dh e^{-\frac{(h - \beta \Gamma_1^2 s_A)^2}{2\Gamma_1^2}} P_h(A \rightarrow B) \quad (\text{C.4})$$

where $P_h(A \rightarrow B)$ is the transition probability given h . We assume that $P_h(A \rightarrow B)$ satisfies detailed balance

$$\frac{P_h(A \rightarrow B)}{P_h(B \rightarrow A)} = e^{-\beta(V(s_B) - V(s_A) - h(s_B - s_A))} \quad (\text{C.5})$$

and we then prove that this implies that the transition probability over a time interval τ , $P(A \rightarrow B, \tau)$, does too. Inserting Eq. (C.5) into Eq. (C.4) we have:

$$\begin{aligned} P(A \rightarrow B) &= \frac{\Gamma_1}{\sqrt{2\pi}} \int dh e^{-\frac{(h - \beta \Gamma_1^2 s_A)^2}{2\Gamma_1^2}} P_h(B \rightarrow A) \\ &\quad \times e^{-\beta(V(s_B) - V(s_A) - h(s_B - s_A))} \\ &= e^{-\beta(V(s_B) - V(s_A))} P(B \rightarrow A) \end{aligned} \quad (\text{C.6})$$

i.e. detailed balance.

-
- 1) L. F. Cugliandolo and J. Kurchan, *Physica A* **263**, 242 (1999).
 - 2) R. Kubo, M. Toda and N. Hashitume, *Statistical Physics II. Nonequilibrium Statistical Mechanics*, Springer-Verlag, 1992.
 - 3) A. J. Bray, *Adv. Phys.* **43**, 357 (1994). D. S. Fisher, cond-mat/9711179 in *Statistical Physics Summer School Lectures on "Fundamental Problems in Statistical Mechanics IX"*, held in Altenberg, Germany, August 20-23, 1997.
 - 4) L. C. E. Struick, *Physical aging in amorphous systems and other materials* (Elsevier, Houston, 1978). G. B. Mc Kenna, *J. Res. NIST* **99**, 169 (1994).
 - 5) L. Lundgren, P. Svedlindh, P. Nordblad and O. Beckmann, *Phys. Rev. Lett.* **51**, 911 (1983). E. Vincent, J. Hammann, M. Ocio, J.-P. Bouchaud and L. F. Cugliandolo; *Slow dynamics and aging in Sitges Conference on Glassy Systems*, M. Rubí ed. (Springer-Verlag, 1997), cond-mat/9607224.
 - 6) F. Alberici-Kious, J.-P. Bouchaud, L. F. Cugliandolo, P. Doussineau and A. Levelut; *Phys. Rev. Lett.* **81**, 4987 (1998).
 - 7) R. L. Leheny and S. Nagel; *Phys. Rev.* **B57**, 5154 (1998).
 - 8) D. Bonn, H. Tanaka, H. Kellay, G. Wegdam and J. Meunier; *Europhys. Lett.* (1998). D. Bonn, H. Tanaka and J. Meunier, *Europhys. Lett.* **45**, 52 (1999).
 - 9) W. van Meegen, T. Mortensen, S. R. Williams and J. Müller, *Phys. Rev.* **E58**, 6073 (1998).
 - 10) J.-P. Bouchaud, L. F. Cugliandolo, J. Kurchan and M. Mézard, *Out of equilibrium dynamics in spin-glasses and other glassy*

- systems*, cond-mat/9702070, in *Spin-glasses and random fields*, A. P. Young ed. (World Scientific, Singapore).
- 11) J-P Bouchaud, cond-mat/9910387.
 - 12) A. Crisanti and H. Sompolinsky; Phys. Rev. **A36**, 4922 (1987). H. Horner, Z. Phys. **B57** (1984) 29; *ibid.* (1984) 39.
 - 13) L. F. Cugliandolo, J. Kurchan, P. Le Doussal and L. Peliti; Phys. Rev. Lett. **78**, 350 (1997).
 - 14) A. Onuki, J. Phys. **9**, 6119 (1997). A. J. Wagner and J. M. Yeomans, cond-mat/9904033. N. P. Rapapa and A. J. Bray, cond-mat/9904396.
 - 15) J. Kurchan, *Rheology, and how to stop aging*, cond-mat/9812347. Proceedings of *Jamming and Rheology: constrained dynamics on microscopic and macroscopic scales*, ITP, Santa Barbara, 1997.
 - 16) L. Berthier, J-L Barrat and J. Kurchan, cond-mat/9910305.
 - 17) G. Chaveteau, R. Tabary, M. Renard and A. Omari, SPE 50752 in *SPE International symposium in oilfield chemistry*, Houston, Texas (1999).
 - 18) M. Jaeger, S. R. Nagel and R. P. Behringer, Rev. Mod. Phys. **68**, 1259 (1996).
 - 19) L.F. Cugliandolo, J. Kurchan and L. Peliti, Phys. Rev. **E55**, 3898 (1997).
 - 20) P. Hanggi, P. Talkner and M. Borkovec, Rev. Mod. Phys. **62**, 251 (1990).
 - 21) C. de Dominicis and P. C. Martin, J. Math. Phys. **5**, 14 (1964), *ibid* 31 (1964). L. Dolan and R. Jackiw, Phys. Rev. **D9**, 2904 (1974), *ibid* 3320 (1974). J. M. Cornwall, R. Jackiw and E. Tomboulis, Phys. Rev. **D10**, 2428 (1974).
 - 22) K. Binder and A. P. Young; Rev. Mod. Phys. **58**, 801 (1986). M. Mézard, G. Parisi and M. A. Virasoro; *Spin glass theory and beyond* (World Scientific, Singapore, 1987). K. H. Fischer and J. A. Hertz, *Spin Glasses*, (Cambridge Univ. Press, 1991).
 - 23) L. F. Cugliandolo and G. Lozano, Phys. Rev. Lett. **80**, 4979 (1998) and Phys. Rev. **B59**, 915 (1999).
 - 24) T. Giamarchi and P. Le Doussal, *Statics and Dynamics of Disordered Elastic Systems*, cond-mat/9705096, in *Spin Glasses and Random Fields*, ed. A.P. Young, World Scientific (Singapore) 1998.
 - 25) A. Georges, G. Kotliar, W. Krauth and M. Rozenberg; Rev. Mod. Phys. **68**, 13 (1996).
 - 26) W. Götze, in *Liquids, freezing and glass transition*, JP Hansen, D. Levesque, J. Zinn-Justin eds. (North Holland, 1989). W. Götze and L. Sjögren, Rep. Prog. Phys. **55**, 241 (1992). W. Götze, J. Phys. **11**, 1 (1999).
 - 27) L. F. Cugliandolo and J. Kurchan, Phys. Rev. Lett. **71**, 173 (1993); J. Phys. **A27**, 5749 (1994); Phil. Mag. **B71**, 501 (1995).
 - 28) T. Natterman, *Theory of the Random Field Ising Model in Spin-Glasses and Random Fields*, ed. A. P. Young, World Scientific (Singapore) 1998.
 - 29) J. Kurchan and L. Laloux, J. Phys. **A29**, 1929 (1996).
 - 30) G. Biroli, cond-mat/9909415.
 - 31) L. F. Cugliandolo, D. S. Dean and J. Kurchan, Phys. Rev. Lett. **79**, 2168 (1997).
 - 32) P. C. Hohenberg and B. I. Shraiman, Physica **D37**, 109 (1989).
 - 33) A phenomenological definition of ‘fictive temperatures’ has been intensively used to fit experimental data of glasses. We refer the reader to Ref.¹⁹⁾ for a comparison of the fictive temperatures to T^{EFF} .
 - 34) For another (inequivalent) definition of temperatures out of equilibrium based on the ‘inherent structure’ ideas see W. Kob, F. Sciortino and P. Tartaglia, cond-mat/9905090. F. Sciortino, W. Kob and P. Tartaglia, cond-mat/9906081.
 - 35) Th. M. Nieuwenhuizen, Phys. Rev. Lett. **80**, 5581 (1998) and cond-mat/9807161.
 - 36) Let us stress that we call ‘temperature’ a quantity that is directly measurable and is related to the zero-th law of thermodynamics.
 - 37) Several aspects of the measurement were clarified by J-L Barrat, private communication.
 - 38) The importance of working with a thermometer that responds in a single time-scale is discussed in R. Exartier and L. Peliti, cond-mat/99010412.
 - 39) N. E. Israeloff and T. S. Grigera, Europhys. Lett. **43**, 308 (1998). T. S. Grigera and N. E. Israeloff, cond-mat/9904351.
 - 40) L. F. Cugliandolo, D. R. Grempel, J. Kurchan and E. Vincent, cond-mat/9905330.
 - 41) A. Barrat, Phys. Rev. **E57**, 3629 (1998). D. Stariolo and S. Cannas, cond-mat/9903136. G. Parisi, F. Ricci-Tersenghi and J. Ruiz-Lorenzo, Phys. Rev. **B57**, 13617 (1998). L. Berthier, J-L Barrat and J. Kurchan, cond-mat/9903091.
 - 42) J-O Andersson, J. Mattson and P. Svedlindh, Phys. Rev. **B46**, 8297 (1992). S. Franz and H. Rieger, J. Stat. Phys. **79**, 749 (1995). E. Marinari, G. Parisi, F. Ricci-Tersenghi, J. J. Ruiz-Lorenzo, J. Phys. **A31**, 2611 (1998). H. Yoshino, K. Hukushima, K. H. Takayama, Prog. Theor. Phys. Supp. 107 (1997). E. Marinari, G. Parisi F. Ricci-Tersenghi and J. Ruiz-Lorenzo, cond-mat/9910232.
 - 43) G. Parisi; Phys. Rev. Lett. **79**, 3660 (1997). W. Kob and J-L Barrat, Physica **A263**, 234 (1999). J-L Barrat and W. Kob, J. Phys. **11** A247 (1999). W. Kob and J-L Barrat, cond-mat/9905248.
 - 44) H. Yoshino, Phys. Rev. Lett. **81**, 1493 (1998) and in preparation.
 - 45) G. Parisi and M. A. Virasoro, J. Phys. (France) **50**, 3317 (1986).
 - 46) T. R. Kirkpatrick and D. Thirumalai, Phys. Rev. **36**, 5388 (1987).
 - 47) R. Monasson, Phys. Rev. Lett. **75**, 2847 (1995).
 - 48) A. E. Allahverdyan and B. Saakian, Phys. Rev. **E58**, 5201 (1998). A. E. Allahverdyan and Th. M. Nieuwenhuizen, cond-mat/9907143. A. E. Allahverdyan, Th. M. Nieuwenhuizen and B. Saakian, cond-mat/9907090.
 - 49) S. Franz and M. A. Virasoro, cond-mat/9907483.
 - 50) A similar very rich example, in spite of its simplicity, has been studied in R. Exartier and L. Peliti, Phys. Lett. **A261** 94 (1999).
 - 51) The process being non-Markovian, one has to specify not only the position x' at a previous time, but also the distribution of all previous positions. Here we consider the stationary measure: $P(x, t; x', t_w)$ stands for the joint probability of $(x, t; x', t_w)$ integrated over all histories from $-\infty$ to t .
 - 52) H. Sompolinsky and A. Zippelius, Phys. Rev. Lett. **47**, 359 (1981), Phys. Rev. **B25**, 6860 (1982). H. Sompolinsky, Phys. Rev. Lett. **47**, 935 (1981), Phil. Mag. **50**, 285 (1984). H. Eissfeller and M. Opper, Phys. Rev. Lett. **68**, 2094 (1992). S. Franz and M. Mézard; Europhys. Lett. **26**, 209 (1994); Physica **A209**, 48 (1994).
 - 53) J-P Hansen and I. R. Mc Donald; *Theory of simple liquids* (Academic Press, London, 1976).
 - 54) ERRATUM: the boson-fermion transformation in Ref.¹⁾ should have read $K(t_a) \rightarrow K(t_s) + \epsilon \bar{\theta}_a$, $\bar{\theta}_a \rightarrow \bar{\theta}_a + \bar{\epsilon}$, $\dot{K}(t_a)\theta_a \rightarrow \dot{K}(t_a)\theta_a + \epsilon T$.
 - 55) Note that we use the set of auxiliary variables y_i to represent the original variable h , unlike the usual situation in which auxiliary variables represent the bath.